arXiv:0912.1138v2 [cond-mat.stat-mech] 30 Jun 2011

The Correlation Functions of the XXZ Heisenberg Chain for Zero or Infinite Anisotropy and Random Walks of Vicious Walkers[*][†]

N. M. Bogoliubov*, **C. Malyshev**$^\diamond$

Steklov Mathematical Institute, St.-Petersburg Department, RAS
Fontanka 27, St.-Petersburg, 191023, Russia
* e-mail: *bogoliub@pdmi.ras.ru*
$^\diamond$ e-mail: *malyshev@pdmi.ras.ru*

Abstract

The XXZ Heisenberg chain is considered for two specific limits of the anisotropy parameter: $\Delta \to 0$ and $\Delta \to -\infty$. The corresponding wave functions are expressed by means of the symmetric Schur functions. Certain expectation values and thermal correlation functions of the ferromagnetic string operators are calculated over the base of N-particle Bethe states. The thermal correlator of the ferromagnetic string is expressed through the generating function of the lattice paths of random walks of vicious walkers. A relationship between the expectation values obtained and the generating functions of strict plane partitions in a box is discussed. Asymptotic estimate of the thermal correlator of the ferromagnetic string is obtained in the limit of zero temperature. It is shown that its amplitude is related to the number of plane partitions.

Keywords: XXZ Heisenberg chain, Schur functions, random walks, plane partitions

---

[*] Extended talk at the Conference "Conformal Field Theory, Integrable Systems, and Liouville Gravity" (Chernogolovka, June 30 – July 2, 2009)

[†] Partially supported by RFBR (No. 07-01-00358) and by the Russian Academy of Sciences program 'Mathematical Methods in Non-Linear Dynamics'

1 Introduction

1.1 XXZ Heisenberg chain

A system of spin *1/2* particles occupying sites of one-dimensional lattice, widely known as the quantum *XYZ Heisenberg chain* [1], has attracted considerable attention both in theoretical and mathematical physics, and it has been thoroughly investigated for a long time [2–8]. The *Quantum Inverse Scattering Method*, developed for solving the integrable models of quantum field theory and statistical physics [9, 10], has been also used to investigate the XYZ Heisenberg chain [11, 12]. An important special case of the XYZ model, so-called, XXZ spin chain also attracts considerable attention [13, 14, 18–20]. The Hamiltonian of the XXZ magnet has been diagonalized by the coordinate *Bethe Ansatz Method* in [3–5]. The *Algebraic Bethe Ansatz* has been used in [11] to solve the XXZ model. The problem of calculation of the correlation functions of the model in question in the framework of the Algebraic Bethe Ansatz has required serious efforts [13–17].

The random walks is a classical problem both for combinatorics and statistical physics. The problem of enumeration of the paths made by the, so-called, *vicious* walkers on the one-dimensional lattice has been formulated and investigated in details by Fisher [21]. The problem mentioned still continues to attract considerable attention both of physicists and mathematicians [22–34].

Random walks on one-dimensional periodic lattice can be related to the correlation functions of the XX Heisenberg magnet. Certain operator averages taken over the ferromagnetic state of the XX model play a role of the generating functions of the number of paths traced by the vicious walkers. The problem of enumeration of trajectories of vicious walkers by means of the correlation functions of the XX model has been studied in the series of papers [35–38]. The approach of Refs. [35–38] will be further explored in the present paper. We shall consider two limits of the XXZ spin chain from the point of view of random walks of vicious walkers, as well as from a viewpoint of enumeration of boxed plane partitions.

Let us begin with the XXZ model defined on one-dimensional lattice consisting of $M+1$ sites labeled by elements of the set $\mathcal{M} \equiv \{0 \le k \le M,\, k \in \mathbb{Z}\}$, $M+1 = 0 \pmod 2$. The corresponding spin Hamiltonian is defined, in absence of external magnetic field, as follows:

$$\hat{H}_{XXZ} = -\frac{1}{2}\sum_{k=0}^{M}(\sigma^-_{k+1}\sigma^+_k + \sigma^+_{k+1}\sigma^-_k + \frac{\Delta}{2}\,(\sigma^z_{k+1}\sigma^z_k - 1))\,, \tag{1}$$

where the parameter $\Delta \in \mathbb{R}$ describes the internal anisotropy of the model. For instance, the choice $\Delta = \pm 1$ corresponds to, so-called, isotropic XXX spin chain solved in [2]. The local spin operators $\sigma^\pm_k = \frac{1}{2}(\sigma^x_k \pm i\sigma^y_k)$ and σ^z_k, dependent on the lattice argument $k \in \mathcal{M}$, are defined as $(M+1)$-fold tensor products as follows:

$$\sigma^\#_k = \sigma^0 \otimes \cdots \otimes \sigma^0 \otimes \underbrace{\sigma^\#}_{k} \otimes \sigma^0 \otimes \cdots \otimes \sigma^0\,, \tag{2}$$

where σ^0 is 2×2 unit matrix, and $\sigma^\#$ at kth place denotes a Pauli matrix, $\sigma^\# \in \mathfrak{su}(2)$ (superscript # implies either x, y, z or $\pm$). Therefore, the spin operators act over the state-space $\mathfrak{H}_{M+1}$ given by the tensor product of $M+1$ copies of the linear spaces $\mathfrak{h}_k \equiv \mathbb{C}^2$: $\mathfrak{H}_{M+1} = \bigotimes\limits_{k=0}^{M} \mathfrak{h}_k$. The commutation rules for the spin operators are given by the relations:

$$[\,\sigma^+_k, \sigma^-_l\,] \,=\, \delta_{k,l}\,\sigma^z_l\,, \quad [\,\sigma^z_k, \sigma^\pm_l\,] \,=\, \pm 2\,\delta_{k,l}\,\sigma^\pm_l\,.$$

The linear space $\mathbb{C}^2$ is spanned over the spin "up" and "down" states ($|\uparrow\rangle$ and $|\downarrow\rangle$, respectively) providing a natural basis so that

$$|\uparrow\rangle \equiv \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \qquad |\downarrow\rangle \equiv \begin{pmatrix} 0 \\ 1 \end{pmatrix}.$$

The space $\mathfrak{H}_{M+1}$ is spanned over of the state-vectors $\bigotimes\limits_{k=0}^{M} |s\rangle_k$, where s implies either $\uparrow$ or $\downarrow$. The periodic boundary conditions $\sigma^{\#}_{k+(M+1)} = \sigma^{\#}_k$ are imposed.

To represent N-particle state-vectors of the model, $|\,\Psi_N(u_1, \ldots, u_N)\rangle$, let the sites with spin "down" states be labeled by the coordinates μ_i, $1 \le i \le N$. These coordinates form a strict partition $\boldsymbol{\mu} \equiv (\mu_1, \mu_2, \ldots, \mu_N)$, where $M \ge \mu_1 > \mu_2 > \ldots > \mu_N \ge 0$. There is a correspondence between each partition and an appropriate sequence of zeros and unities of the form: $\big\{e_k \equiv e_k(\boldsymbol{\mu})\big\}_{k\in\mathcal{M}}$, where $e_k \equiv \delta_{k,\mu_n}$, $1 \le n \le N$. This correspondence enables another convenient expression for the strict partitions: $\boldsymbol{\mu} = (M^{e_M}, \ldots, 1^{e_1}, 0^{e_0})$. It is meant here that any non-negative integer $k \in \mathcal{M}$ appears e_k times in the present configuration, and the condition $\sum\limits_{k=0}^{M} e_k = N$ is respected. The Hamiltonian (1) is diagonalized *via* the ansatz:

$$|\,\Psi_N(\mathbf{u})\rangle \;=\; \sum_{\{e_k(\boldsymbol{\mu})\}_{k\in\mathcal{M}}} \chi^{XXZ}_{\boldsymbol{\mu}}(\mathbf{u}) \prod_{k=0}^{M} (\sigma^-_k)^{e_k}\, |\Uparrow\rangle\,, \tag{3}$$

where the sum is taken over C^N_M strict partitions $\boldsymbol{\mu}$. The state $|\Uparrow\rangle$ in (3) is the fully polarized state with all spins "up": $|\Uparrow\rangle \equiv \bigotimes\limits_{n=0}^{M} |\uparrow\rangle_n$. Besides, it is proposed to use bold-faced letters as short-hand notations for appropriate N-tuples of numbers: for instance, $\mathbf{u}$ instead of $(u_1, \ldots, u_N)$, etc. Therefore, the wave function $\chi^{XXZ}_{\boldsymbol{\mu}}(\mathbf{u})$ in (3) is of the form:

$$\chi^{XXZ}_{\boldsymbol{\mu}}(\mathbf{u}) \;=\; \sum_{S_{p_1,p_2,\ldots,p_N}} \mathcal{A}_S(\mathbf{u})\, u^{2\mu_1}_{p_1} u^{2\mu_2}_{p_2} \ldots u^{2\mu_N}_{p_N}\,, \tag{4}$$

where summation goes over all elements of the symmetric group of permutations $S_{p_1,p_2,\ldots,p_N} \equiv S\begin{pmatrix} 1, & 2, & \ldots, & N \\ p_1, & p_2, & \ldots, & p_N \end{pmatrix}$. The amplitude $\mathcal{A}_S$ is given by the product:

$$\mathcal{A}_S(\mathbf{u}) \;\equiv\; \prod_{1\le j<i\le N} \frac{1 - 2\Delta u^2_{p_i} + u^2_{p_i} u^2_{p_j}}{u^2_{p_i} - u^2_{p_j}}\,. \tag{5}$$

The state-vectors (3) are the eigen-states of the Hamiltonian (1),

$$\hat{H}_{XXZ}\, |\Psi_N(\mathbf{u})\rangle \;=\; E_N(\mathbf{u})\, |\Psi_N(\mathbf{u})\rangle\,,$$

if and only if the variables u_l ($1 \le l \le N$) satisfy the *Bethe equations*

$$u_l^{2(M+1)} = (-1)^{N-1} \prod_{k=1}^{N} \frac{1 - 2\Delta u^2_l + u^2_l u^2_k}{1 - 2\Delta u^2_k + u^2_l u^2_k}\,. \tag{6}$$

The corresponding eigen-energies are given by

$$E_N(\mathbf{u}) \;=\; -\frac{1}{2} \sum_{i=1}^{N} (u^2_i + u^{-2}_i - 2\Delta)\,. \tag{7}$$

1.2 Outline of the limiting models and the problem

In our paper we shall consider two special cases of the XXZ model, namely the $\Delta \to 0$ and $\Delta \to -\infty$ limits. The $\Delta \to 0$ limit known as the XX Heisenberg chain is a most popular and studied one [39–47]. The Hamiltonian of the XX model describes the nearest-neighbor interactions of spin "up" $|\uparrow\rangle$ and spin "down" $|\downarrow\rangle$ states located on sites of the periodic chain in zero magnetic field as follows:

$$\hat{H}_{XX} \equiv -\frac{1}{2}\sum_{k=0}^{M}(\sigma_{k+1}^{-}\sigma_{k}^{+} + \sigma_{k+1}^{+}\sigma_{k}^{-})\,. \tag{8}$$

It is crucial that the system described by the Hamiltonian (8) is equivalent to free fermions [41]. The case of the XX magnet can also be deduced by taking the limit of infinite on-site repulsion in the boson Hubbard model (the "hard-core" bosons) [48]. Therefore, the XX model is interesting for investigating the (quantum) phase diagram of the Hubbard model, as well as for description of the Frenkel excitons [49]. In the last few years, the XX model has attracted attention in connection with the quantum information and computation theory [50, 51].

The state-vector of the XX model and the correspondent Bethe equations are obtained at $\Delta = 0$ from (4) and (6), respectively. Up to an irrelevant pre-factor, the wave functions of the model are equal to

$$\chi_{\boldsymbol{\mu}}^{XX}(\mathbf{u}) = \det(u_j^{2\mu_k})_{1\le j,k\le N} \prod_{1\le n<l\le N} (u_l^2 - u_n^2)^{-1}, \tag{9}$$

and the Bethe equations are [41]:

$$u_j^{2(M+1)} = (-1)^{N-1}, \quad 1 \le j \le N. \tag{10}$$

The substitution $u_j^2 = e^{i\theta_j}$ brings these equations to the exponential form,

$$e^{i(M+1)\theta_j} = (-1)^{N-1}, \tag{11}$$

with the solutions:

$$\theta_j = \frac{2\pi}{M+1}\Big(I_j - \frac{N-1}{2}\Big), \tag{12}$$

where I_j are integers or half-integers depending on whether N is odd or even. It is sufficient to consider a set of N different numbers I_j satisfying the condition: $M \ge I_1 > I_2 > \cdots > I_N \ge 0$. The notation $\boldsymbol{\theta}$ for N-tuple $(\theta_1, \theta_2, \ldots, \theta_N)$ of solutions (12) will be especially convenient for usage below in order to stress that one is concerned with the solution of the Bethe equation. Otherwise, it is appropriate to use $\mathbf{u}$ as an indication that arbitrary set of parameters is meant. It follows from (7) that the eigen-energies of the XX model are equal to

$$E_N^{XX}(\boldsymbol{\theta}) \;=\; -\sum_{j=1}^{N}\cos\theta_j \;=\; -\sum_{j=1}^{N}\cos\left(\frac{2\pi}{M+1}\Big(I_j - \frac{N-1}{2}\Big)\right). \tag{13}$$

The ground state of the model corresponds to the following solutions of the Bethe equations:

$$\theta_j = \frac{2\pi}{M+1}\Big(N - j - \frac{N-1}{2}\Big), \quad 1 \le j \le N. \tag{14}$$

Less studied limit of the XXZ model is the *Strong Anisotropy* (SA) limit $\Delta \to -\infty$ [4,8,52–55]. In this limit the behavior of the system is described by the effective Hamiltonian which is formally equivalent to the XX Hamiltonian supplied with requirement forbidding two spin "down" states to occupy any pair of nearest-neighboring sites [52, 53]:

$$\hat{H}_{\mathrm{SA}} = -\frac{1}{2} \sum_{k=0}^{M} \mathcal{P}(\sigma_{k+1}^{-}\sigma_{k}^{+} + \sigma_{k+1}^{+}\sigma_{k}^{-})\mathcal{P}\,, \tag{15}$$

where the projector $\mathcal{P}$ cuts out the states with the spin "down" states at any pair of nearest-neighboring sites: $\mathcal{P} \equiv \prod_{k=0}^{M}(1 - \hat{q}_{k+1}\hat{q}_k)$. The local projectors onto the spin "up" and "down" states are equal to:

$$\check{q}_k \,\equiv\, \frac{1}{2}\,(\sigma_k^0 + \sigma_k^z)\,, \quad \hat{q}_k \,\equiv\, \frac{1}{2}\,(\sigma_k^0 - \sigma_k^z)\,, \quad \check{q}_k + \hat{q}_k \,=\, \mathbb{I}, \qquad k \in \mathcal{M}\,, \tag{16}$$

where the operators $\sigma_k^{\#}$ are defined by (2).

In the limit $\Delta \to -\infty$, the wave function (4) takes the form:

$$\chi_{\boldsymbol{\mu}}^{\mathrm{SA}}(\mathbf{u}) = \det(u_j^{2(\mu_k - N + k)})_{1\le j,k\le N} \prod_{1\le n<l\le N} (u_l^2 - u_n^2)^{-1}\,, \tag{17}$$

where the coordinates of the spin "down" states form strict decreasing partition $\boldsymbol{\mu}$ (as in (4) and (9)), i.e., $M \ge \mu_1 > \mu_2 > \ldots > \mu_N \ge 0$. It follows from (17) that the wave function is not equal to zero if and only if the elements μ_i, $1 \le i \le N$, satisfy the *exclusion condition*: $\mu_i > \mu_{i+1} + 1$. It is crucial that in the considered limit the occupation of nearest sites is forbidden, and the *hard-core diameter* equal to duplicated inter-site separation arises. The Bethe equations of the model take the form:

$$e^{i(M+1-N)\theta_k} = (-1)^{N-1} \prod_{j=1}^{N} e^{-i\theta_j}\,, \quad 1 \le k \le N\,, \tag{18}$$

and have the solutions

$$\theta_j = \frac{2\pi}{M+1-N}\Bigl(I_j - \frac{N-1}{2} - P\Bigr), \tag{19}$$

where $P \equiv \frac{1}{2\pi}\sum_{j=1}^{N}\theta_j$, and I_j are integers or half-integers depending on N being odd or even, satisfying the condition $M - N \ge I_1 > I_2 > \cdots > I_N \ge 0$. The ground state of the model is defined by the solutions

$$\theta_j = \frac{2\pi}{M+1-N}\Bigl(N - j - \frac{N-1}{2}\Bigr), \tag{20}$$

while $P = 0$ is fulfilled. The eigen-energy of the model is:

$$E_N(\boldsymbol{\theta}) \,=\, -\sum_{j=1}^{N} \cos\left(\frac{2\pi}{M+1-N}\Bigl(I_j - \frac{N-1}{2} - P\Bigr)\right)\,. \tag{21}$$

Similarity between these two limits, (8) and (15), is that their wave functions are expressible through the *Schur functions* [56]:

$$\begin{aligned} S_{\boldsymbol{\lambda}}(\mathbf{x}) \,\equiv\, S_{\boldsymbol{\lambda}}(x_1, x_2, \dots, x_N) \,&\equiv\, \frac{\det(x_j^{\lambda_k+N-k})_{1\le j,k\le N}}{\det(x_j^{N-k})_{1\le j,k\le N}} \\ &=\, \det(x_j^{\lambda_k+N-k})_{1\le j,k\le N} \prod_{1\le n<l\le N} (x_l - x_n)^{-1}\,, \end{aligned} \tag{22}$$

where $\boldsymbol{\lambda}$ denotes the partition $(\lambda_1, \lambda_2, \dots, \lambda_N)$ being N-tuple of non-increasing non-negative integers: $L \ge \lambda_1 \ge \lambda_2 \ge \, \dots \, \ge \lambda_N \ge 0$. Indeed, any strict partition $M \ge \mu_1 > \mu_2 > \, \dots \, > \mu_N \ge 0$ and non-strict partition $M + 1 - N \ge \lambda_1 \ge \lambda_2 \ge \cdots \ge \lambda_N \ge 0$ (denoted as $\boldsymbol{\mu}$ and $\boldsymbol{\lambda}$, respectively) can be related by means of the relation $\lambda_j = \mu_j - N + j$, where $1 \le j \le N$. In other terms, $\boldsymbol{\lambda} = \boldsymbol{\mu} - \boldsymbol{\delta}$, where $\boldsymbol{\delta}$ is the strict partition $(N-1, N-2, \dots, 1, 0)$. So the wave function of the XX model (9) may be represented as

$$\chi^{XX}_{\boldsymbol{\mu}}(\mathbf{u}) = S_{\boldsymbol{\lambda}}(\mathbf{u}^2)\,. \tag{23}$$

Any strict partition $\boldsymbol{\mu}$ with the elements respecting the exclusion condition $\mu_i > \mu_{i+1} + 1$ is connected with the non-strict partition $\widetilde{\boldsymbol{\lambda}}$ by the relation $\widetilde{\boldsymbol{\lambda}} = \boldsymbol{\mu} - 2\boldsymbol{\delta}$, where $M + 2(1-N) \ge \widetilde{\lambda}_1 \ge \widetilde{\lambda}_2 \ge \cdots \ge \widetilde{\lambda}_N \ge 0$. Therefore, Eq. (17) is re-expressed:

$$\chi^{\mathrm{SA}}_{\boldsymbol{\mu}}(\mathbf{u}) = S_{\widetilde{\boldsymbol{\lambda}}}(\mathbf{u}^2)\,. \tag{24}$$

It is useful to remind a graphical picture (see Figure 1) of the correspondence between strict partitions $\boldsymbol{\mu}$ and non-strict partitions $\boldsymbol{\lambda}$ used in (23). Namely, to each partition $\boldsymbol{\lambda}$ we associate a set of N-tuples $\mathfrak{G}(\boldsymbol{\lambda})$ as follows [60]:

$$\mathfrak{G}(\boldsymbol{\lambda}) \,=\, \Big\{ \lambda_j - j + \frac{1}{2} \,\Big|\, 1 \le j \le N \Big\} \subset \mathbb{Z} + \frac{1}{2}\,.$$

On another hand, any non-strict partition $\boldsymbol{\lambda}$ can be represented as a rectangular table (the Young table) consisting of N columns so that λ_i, $\forall i$, is the height of i^{th} column ($\lambda_i \le M - N + 1$). We shift each element of the set $\mathfrak{G}(\boldsymbol{\lambda})$ by $N + \frac{1}{2}$. Then we assign the numbers thus obtained to the projections along the vertical dashed lines onto the horizontal axis. The set of points on the horizontal axis just provides the strict partition $\boldsymbol{\mu}$. For instance, the diagram on Figure 1 is drown for $M = 8$ and $N = 4$, and we have got respectively: $\boldsymbol{\lambda} = (5, 3, 2, 2)$ and $\boldsymbol{\mu} = (8, 5, 3, 2)$.

In our paper we shall study the thermal correlation function of the states (would be called as *ferromagnetic strings*) with no spins down on the last $n+1$ sites of the lattice. We are going to consider the expectation value defined by the ratio:

$$\mathcal{T}(\boldsymbol{\theta}, n, \beta) \,\equiv\, \frac{\langle \Psi_N(\boldsymbol{\theta}) \mid \bar{\Pi}_n \, e^{-\beta \hat{H}} \, \bar{\Pi}_n \mid \Psi_N(\boldsymbol{\theta}) \rangle}{\langle \Psi_N(\boldsymbol{\theta}) \mid \Psi_N(\boldsymbol{\theta}) \rangle}\,, \qquad \bar{\Pi}_n \equiv \prod_{j=M-n}^{M} \check{q}_j\,, \tag{25}$$

where $\beta \in \mathbb{C}$, and the projector $\bar{\Pi}_n$ is expressed by means of $\check{q}_j$ (16). Besides, $\hat{H}$ in $\mathcal{T}(\boldsymbol{\theta}, n, \beta)$ (25) implies either $\hat{H}_{XX}$ (8) or $\hat{H}_{\mathrm{SA}}$ (15), and $\boldsymbol{\theta}$ indicates that the eigen-state $\mid \Psi_N(\boldsymbol{\theta})\rangle$ is calculated for solution of the Bethe equation (11) or (18), respectively. Our calculations will



Figure 1: The strict partition $(\mu_1, \mu_2, \mu_3, \mu_4)$ and the corresponding Young table.

be based on the similarity of the problem under consideration with the problem of enumeration of admissible lattice paths made by random vicious walkers. We shall extensively use the relation for the Schur functions (22) which is due to the Binet–Cauchy formula [57]:

$$\begin{aligned} & \sum_{\boldsymbol{\lambda}\subseteq\{L^N\}} S_{\boldsymbol{\lambda}}(x_1^2,\dots,x_N^2) S_{\boldsymbol{\lambda}}(y_1^2,\dots,y_N^2) \\ = \; & \det(T_{jk})_{1\le j,k\le N} \prod_{1\le k<j\le N} \left(y_j^2 - y_k^2\right)^{-1} \prod_{1\le m<l\le N} \left(x_l^2 - x_m^2\right)^{-1} . \end{aligned} \tag{26}$$

The entries of the matrix T_{jk} take the form:

$$T_{jk} = \frac{1-(x_k y_j)^{2(N+L)}}{1-(x_k y_j)^2} . \tag{27}$$

Summation in (26) goes over all non-strict partitions $\boldsymbol{\lambda}$ into at most N parts so that each is less than L: $L \ge \lambda_1 \ge \lambda_2 \ge \cdots \ge \lambda_N \ge 0$. The notation for the range of summation in (26) will be extensively used in the rest of the paper.

2 XX Heisenberg chain

2.1 The Bethe states and form-factors

Before to proceed with calculation of $\mathcal{T}(\boldsymbol{\theta}, n, \beta)$ (25), let us apply our approach to more familiar examples. First of all, let us specialize, with regard at (23), the state-vector (3) and its conjugated as follows:

$$\begin{aligned} |\,\Psi_N(\mathbf{u})\rangle &= \sum_{\boldsymbol{\lambda}\subseteq\{(M+1-N)^N\}} S_{\boldsymbol{\lambda}}(\mathbf{u}^2) \prod_{k=0}^{M} (\sigma_k^-)^{e_k}\, |\Uparrow\rangle , \\ \langle \Psi_N(\mathbf{v})\,| &= \sum_{\boldsymbol{\lambda}\subseteq\{(M+1-N)^N\}} \langle\Uparrow| \prod_{k=0}^{M} (\sigma_k^+)^{\tilde{e}_k} S_{\boldsymbol{\lambda}}(\mathbf{v}^{-2}) , \end{aligned} \tag{28}$$

where summation goes over all non-strict partitions $\boldsymbol{\lambda}$, which are related to the non-strict partitions $\boldsymbol{\mu} = \boldsymbol{\lambda} + \boldsymbol{\delta}$. Because of the orthogonality relation

$$\langle \Uparrow | \prod_{k=0}^{M} (\sigma_k^+)^{\tilde{e}_k} \prod_{l=0}^{M} (\sigma_l^-)^{e_l} | \Uparrow \rangle = \prod_{n=0}^{M} \delta_{\tilde{e}_n e_n} ,$$

the scalar product of the state-vectors (28) takes the form:

$$\langle \Psi_N(\mathbf{v}) \mid \Psi_N(\mathbf{u}) \rangle \;=\; \sum_{\boldsymbol{\lambda} \subseteq \{(M+1-N)^N\}} S_{\boldsymbol{\lambda}}(\mathbf{v}^{-2}) S_{\boldsymbol{\lambda}}(\mathbf{u}^2) \;=\; \frac{\det(T_{kj})_{1\le k,j\le N}}{\mathcal{V}(\mathbf{u}^2)\mathcal{V}(\mathbf{v}^{-2})} . \tag{29}$$

We use in (29) the notation for the Vandermonde determinant,

$$\mathcal{V}(\mathbf{u}^2) \;\equiv\; \prod_{1\le m<l\le N} (u_l^2 - u_m^2) , \tag{30}$$

while the entries T_{kj} of the matrix of the size $N \times N$ are of the form:

$$T_{kj} = \frac{1 - (u_k^2/v_j^2)^{M+1}}{1 - u_k^2/v_j^2} . \tag{31}$$

Equations (29) and (31) are specifications of Eqs. (26) and (27), respectively. On the solutions (12), the entries (31) are equal to $T_{jk} = (M+1)\delta_{jk}$, where l'Hospital rule is taken into account on the principle diagonal. Let us use the exponential parametrization for the solutions of the Bethe equations in compact form:

$$\mathbf{u}^2 = e^{i\boldsymbol{\theta}} , \qquad e^{i\boldsymbol{\theta}} \equiv (e^{i\theta_1}, e^{i\theta_2}, \ldots, e^{i\theta_N}) , \tag{32}$$

where the "angle" notation (12) or (19) is meant. Thus, the answer for squared norm $\mathcal{N}^2(\boldsymbol{\theta}) \equiv \langle \Psi_N(\boldsymbol{\theta}) \mid \Psi_N(\boldsymbol{\theta}) \rangle$ of the Bethe eigen-vector (28) arises as follows:

$$\mathcal{N}^2(\boldsymbol{\theta}) \;=\; \frac{(M+1)^N}{\mathcal{V}(e^{i\boldsymbol{\theta}})\mathcal{V}(e^{-i\boldsymbol{\theta}})} = \frac{(M+1)^N}{\prod\limits_{1\le m<l\le N} 2(1 - \cos(\theta_l - \theta_m))} . \tag{33}$$

Let us now turn to the ratio:

$$\mathcal{T}(\mathbf{v}, \mathbf{u}, n) \;\equiv\; \frac{\langle \Psi_N(\mathbf{v}) \mid \bar{\Pi}_n \mid \Psi_N(\mathbf{u}) \rangle}{\mathcal{N}(\mathbf{v})\mathcal{N}(\mathbf{u})} , \tag{34}$$

where the projector $\bar{\Pi}_n$ is defined in (25), and $\mathcal{N}^2(\mathbf{u}) = \langle \Psi_N(\mathbf{u}) \mid \Psi_N(\mathbf{u}) \rangle$ for arbitrary parametrization of the state-vectors. With regard at (28), we calculate:

$$\bar{\Pi}_n \mid \Psi_N(\mathbf{u}) \rangle \;=\; \sum_{\boldsymbol{\lambda} \subseteq \{(M-N-n)^N\}} S_{\boldsymbol{\lambda}}(\mathbf{u}^2) \Big(\prod_{k=M-n}^{M} (\sigma_k^-)^0 \Big) \Big(\prod_{k=0}^{M-n-1} (\sigma_k^-)^{e_k} \Big) | \Uparrow \rangle , \tag{35}$$

where summation goes over non-strict partitions $\boldsymbol{\lambda}$ respecting the condition: $M - N - n \ge \lambda_1 \ge \lambda_2 \ge \ldots \ge \lambda_N \ge 0$. Taking into account (28) and (35), and using the Binet-Cauchy formula, we calculate the nominator of (34):

$$\begin{aligned} \langle \Psi_N(\mathbf{v}) \mid \bar{\Pi}_n \mid \Psi_N(\mathbf{u}) \rangle \;&=\; \sum_{\boldsymbol{\lambda} \subseteq \{(M-N-n)^N\}} S_{\boldsymbol{\lambda}}(\mathbf{v}^{-2}) S_{\boldsymbol{\lambda}}(\mathbf{u}^2) \\ &= \frac{1}{\mathcal{V}(\mathbf{u}^2)\mathcal{V}(\mathbf{v}^{-2})} \det \left(\frac{1 - (u_k^2/v_j^2)^{M-n}}{1 - u_k^2/v_j^2} \right)_{1\le j,k\le N} . \end{aligned} \tag{36}$$

Assume that the sets of the parameters $\mathbf{v}$ and $\mathbf{u}$ in (34) coincide and consist of the solutions (12). Then, expression $\mathcal{T}(\boldsymbol{\theta}, n) \equiv \mathcal{T}(e^{i\boldsymbol{\theta}/2}, e^{i\boldsymbol{\theta}/2}, n)$ is related to, so-called, *Emptiness Formation Probability*, which provides the probability of formation of a string of the empty (i.e., spin "up") states on last $n+1$ sites of the lattice. Eventually, Eq. (36) leads to $\mathcal{T}(\boldsymbol{\theta}, n)$ in the same determinantal form as in [41, 43] (see therein for more Refs.):

$$\mathcal{T}(\boldsymbol{\theta}, n) \;=\; \det\Big(\big(1 - \frac{n+1}{M+1}\big)\delta_{jk} + \frac{1 - e^{i(\theta_j - \theta_k)(n+1)}}{(M+1)(1 - e^{i(\theta_k - \theta_j)})}(1 - \delta_{jk})\Big)_{1\le j,k\le N}\,, \tag{37}$$

where the parameters θ_l, $1 \le l \le N$, correspond to the parametrization (12).

2.2 Thermal correlator of ferromagnetic string and random walks of vicious walkers

Let us turn to calculation of the following expectation value:

$$\mathcal{T}(\mathbf{v}, \mathbf{u}, n, \beta) \;\equiv\; \frac{\langle \Psi_N(\mathbf{v}) \mid \bar{\Pi}_n \, e^{-\beta \hat{H}_{XX}} \, \bar{\Pi}_n \mid \Psi_N(\mathbf{u})\rangle}{\mathcal{N}(\mathbf{v})\mathcal{N}(\mathbf{u})}\,, \tag{38}$$

which is clearly reduced to $\mathcal{T}(\mathbf{v}, \mathbf{u}, n)$ (34) at $\beta = 0$. However, our aim is to obtain $\mathcal{T}(\boldsymbol{\theta}, n, \beta)$ (25) just taking the arguments $\mathbf{u}$ and $\mathbf{v}$ in (38) coinciding with the same solution (12).

Let us use the technique presented above to calculation of the nominator of (38). Using (35) (and its conjugated), one gets:

$$\begin{aligned} &\langle \Psi_N(\mathbf{v}) \mid \bar{\Pi}_n \, e^{-\beta \hat{H}_{XX}} \, \bar{\Pi}_n \mid \Psi_N(\mathbf{u})\rangle \\ &= \sum_{\boldsymbol{\lambda}^L,\, \boldsymbol{\lambda}^R \subseteq \{(M-N-n)^N\}} S_{\boldsymbol{\lambda}^L}(\mathbf{v}^{-2}) S_{\boldsymbol{\lambda}^R}(\mathbf{u}^2) F_{\boldsymbol{\mu}^L;\,\boldsymbol{\mu}^R}(\beta)\,. \end{aligned} \tag{39}$$

Summations in (39) run over non-strict partitions $\boldsymbol{\lambda}^L$ and $\boldsymbol{\lambda}^R$ of the same kind as in (35). Superscripts L and R are only to distinguish two independent summations. The corresponding strict partitions $\boldsymbol{\mu}^L$ and $\boldsymbol{\mu}^R$ are defined as follows: $\boldsymbol{\mu}^{L,R} = \boldsymbol{\lambda}^{L,R} + \boldsymbol{\delta}$, where $\boldsymbol{\delta} \equiv (\delta_1, \delta_2, \ldots, \delta_N)$, $\delta_j = N - j$. The notation $F_{\boldsymbol{\mu}^L;\,\boldsymbol{\mu}^R}(\beta)$ implies the following average:

$$\begin{aligned} F_{\boldsymbol{\mu}^L;\,\boldsymbol{\mu}^R}(\beta) \;&\equiv\; F_{\mu_1^L, \mu_2^L, \ldots, \mu_N^L; \mu_1^R, \mu_2^R, \ldots, \mu_N^R}(\beta) \;= \\ &=\; \langle \Uparrow | \sigma^+_{\mu_1^L} \sigma^+_{\mu_2^L} \ldots \sigma^+_{\mu_N^L} e^{-\beta \hat{H}_{XX}} \sigma^-_{\mu_1^R} \sigma^-_{\mu_2^R} \ldots \sigma^-_{\mu_N^R} | \Uparrow \rangle\,, \end{aligned} \tag{40}$$

which is nothing but $2N$-point correlation function over the ferromagnetic state. It is related to enumeration of admissible trajectories which are traced by N vicious walkers traveling over sites of one-dimensional chain [35–38].

Indeed, let $|P_K(\mu_1^R, \ldots, \mu_N^R \to \mu_1^L, \ldots, \mu_N^L)|$ be a number of trajectories consisting of K links made by N vicious walkers in the random turns model. Here, the initial and final positions of the walkers on the sites are given respectively by elements of the strict decreasing partitions $\mu_1^R > \mu_2^R > \cdots > \mu_N^R$ and $\mu_1^L > \mu_2^L > \cdots > \mu_N^L$. We introduce the notation $\mathcal{D}_\ell^K$ for the operator of differentiation of K^{th} order with respect to ℓ at the point $\ell = 0$ [38]. Then, the application of the operator $\mathcal{D}_{\beta/2}^K$ to the correlator (40) results in the average of the type

$$\langle \Uparrow | \sigma^+_{\mu_1^L} \sigma^+_{\mu_2^L} \ldots \sigma^+_{\mu_N^L} (-2\hat{H}_{XX})^K \sigma^-_{\mu_1^R} \sigma^-_{\mu_2^R} \ldots \sigma^-_{\mu_N^R} | \Uparrow \rangle\,.$$

This average provides the numbers $|P_K(\mu_1^R,\dots,\mu_N^R \to \mu_1^L,\dots,\mu_N^L)|$, as it has been established in [36] with the help of the commutation relation

$$[\hat{H}_{XX}, \sigma^-_{\mu_1^R}\sigma^-_{\mu_2^R}\dots\sigma^-_{\mu_N^R}] = \sum_{k=1}^{N} \sigma^-_{\mu_1^R}\dots\sigma^-_{\mu_{k-1}^R}[\hat{H}_{XX}, \sigma^-_{\mu_k^R}]\sigma^-_{\mu_{k+1}^R}\dots\sigma^-_{\mu_N^R}\,. \tag{41}$$

The condition of non-intersection of trajectories of the walkers is expressed by the vanishing of the correlation function (40) for any pair of coinciding indices μ_k^R or μ_p^L. Thus we conclude that the average (39) turns out to be the generating function of the polynomials dependent on $2N$ variables, $u_1^2, u_2^2, \dots, u_N^2$ and $v_1^{-2}, v_2^{-2}, \dots, v_N^{-2}$, as follows:

$$\begin{aligned} &\mathcal{D}^K_{\beta/2}\left[\langle\Psi_N(\mathbf{v}) \mid \bar{\Pi}_n\, e^{-\beta\hat{H}_{XX}}\, \bar{\Pi}_n \mid \Psi_N(\mathbf{u})\rangle\right] = \\ &= \sum_{\boldsymbol{\lambda}^L,\,\boldsymbol{\lambda}^R\subseteq\{(M-N-n)^N\}} |P_K(\boldsymbol{\mu}^R\to\boldsymbol{\mu}^L)|\, S_{\boldsymbol{\lambda}^L}(\mathbf{v}^{-2})\, S_{\boldsymbol{\lambda}^R}(\mathbf{u}^2) \end{aligned} \tag{42}$$

(remind that $\boldsymbol{\mu}^{L,R} = \boldsymbol{\lambda}^{L,R} + \boldsymbol{\delta}$). As it is shown in [38], the number of trajectories consisting of K links, which are traced by N vicious walkers on an axis, i.e, $|P_K(\boldsymbol{\mu}^R\to\boldsymbol{\mu}^L)|$, is expressed through the number of trajectories of the same "length" K, which are traced by a single walker traveling over sites of N-dimensional lattice of infinite extension.

The correlator (40) respects the following equation:

$$\begin{aligned} &\frac{d}{d\beta}\, F_{\mu_1^L,\mu_2^L,\dots,\mu_N^L;\mu_1^R,\mu_2^R,\dots,\mu_N^R}(\beta) = \\ &= \frac{1}{2}\sum_{k=1}^{N}\big(F_{\mu_1^L,\mu_2^L,\dots,\mu_N^L;\mu_1^R,\mu_2^R,\dots,\mu_k^R+1,\dots,\mu_N^R}(\beta) + F_{\mu_1^L,\mu_2^L,\dots,\mu_N^L;\mu_1^R,\mu_2^R,\dots,\mu_k^R-1,\dots,\mu_N^R}(\beta)\big)\,. \end{aligned} \tag{43}$$

Equation (43) has been considered in [36] for the case of the periodic boundary condition with respect to the lattice argument and with the initial condition:

$$F_{\mu_1^L,\mu_2^L,\dots,\mu_N^L;\mu_1^R,\mu_2^R,\dots,\mu_N^R}(0) = \prod_{k=1}^{N}\delta_{\mu_k^L,\mu_k^R}\,.$$

Solution to (43) can be expressed as the determinant of the matrix $\big(F_{k;\,l}(\beta)\big)_{1\le k,l\le N}$ [35,36]:

$$F_{\mu_1^L,\mu_2^L,\dots,\mu_N^L;\mu_1^R,\mu_2^R,\dots,\mu_N^R}(\beta) = \det\big(F_{\mu_k^L;\mu_l^R}(\beta)\big)_{1\le k,l\le N}\,, \tag{44}$$

where the entries respect the following difference-differential equation:

$$\frac{d}{d\beta}\,F_{k;\,l}(\beta) = \frac{1}{2}\,(F_{k+1;l}(\beta) + F_{k-1;l}(\beta))\,. \tag{45}$$

Similar equation can be also obtained for the fixed index l.

It can be checked that the transition amplitude $\langle\Uparrow|\sigma_k^+ e^{-\beta\hat{H}_{XX}}\sigma_l^-|\Uparrow\rangle$ respects (45) [35–38]. This average can be considered as the generating function of the number of walks with random turns of a single pedestrian traveling between 1^{th} and k^{th} sites of (periodic) chain [36,38]. Solution to (45) can be written as the following sum:

$$F_{k;\,l}(\beta) \equiv \frac{1}{M+1}\sum_{s=0}^{M} e^{\beta\cos\phi_s}\, e^{i\phi_s(k-l)}\,, \tag{46}$$

where the parametrization $\phi_s = \dfrac{2\pi}{M+1}\Big(s - \dfrac{M}{2}\Big)$ is used. The periodicity condition with respect of the lattice argument and the "initial" condition $F_{k;\,l}(0) = \delta_{k,l}$ are imposed.

Using (44) and (46), we re-express (40) through the Schur functions (22) and the Vandermonde determinants (30) as follows [35]:

$$
\begin{aligned}
F_{\boldsymbol{\mu}^L;\,\boldsymbol{\mu}^R}(\beta) \;=\; & \frac{1}{(M+1)^N} \sum_{M\geq k_1>k_2\cdots>k_N\geq 0} e^{\beta\sum\limits_{l=1}^{N}\cos(\phi_{k_l})} \\
& \times \mathcal{V}(e^{i\boldsymbol{\phi}})\mathcal{V}(e^{-i\boldsymbol{\phi}})\, S_{\boldsymbol{\lambda}^L}(e^{i\boldsymbol{\phi}}) S_{\boldsymbol{\lambda}^R}(e^{-i\boldsymbol{\phi}})\,.
\end{aligned}
\tag{47}
$$

The parametrization ϕ_s, $0 \leq s \leq M$, used in (47) is the same as in (46). We continue to use bold-faced letters to denote N-tuples of numbers: for instance, $\boldsymbol{\phi}$ corresponds to $(\phi_{k_1}, \phi_{k_2}, \dots, \phi_{k_N})$. We substitute (47) into (39), and use (36) in order to calculate the sums:

$$
\mathcal{P}(\mathbf{v}^{-2}, e^{i\boldsymbol{\phi}}) \;\equiv\; \sum_{\boldsymbol{\lambda}^L} S_{\boldsymbol{\lambda}^L}(\mathbf{v}^{-2})\, S_{\boldsymbol{\lambda}^L}(e^{i\boldsymbol{\phi}})\,, \quad \mathcal{P}(e^{-i\boldsymbol{\phi}}, \mathbf{u}^2) \;\equiv\; \sum_{\boldsymbol{\lambda}^R} S_{\boldsymbol{\lambda}^R}(e^{-i\boldsymbol{\phi}})\, S_{\boldsymbol{\lambda}^R}(\mathbf{u}^2)\,. \tag{48}
$$

The range of summation in (48) is taken as follows: $\boldsymbol{\lambda}^L$, $\boldsymbol{\lambda}^R \subseteq \{(M-N-n)^N\}$. Then we obtain:

$$
\begin{aligned}
& \langle \Psi_N(\mathbf{v}) \mid \bar{\varPi}_n\, e^{-\beta \hat{H}_{XX}}\, \bar{\varPi}_n \mid \Psi_N(\mathbf{u}) \rangle \;= \\
& = \frac{1}{(M+1)^N\, \mathcal{V}(\mathbf{u}^2)\mathcal{V}(\mathbf{v}^{-2})} \sum_{M\geq k_1>k_2\cdots>k_N\geq 0} e^{\beta\sum\limits_{l=1}^{N}\cos(\phi_{k_l})} \times \\
& \times \det\Big(\frac{1-(e^{i\phi_{k_i}} v_j^{-2})^{M-n}}{1-e^{i\phi_{k_i}} v_j^{-2}}\Big)_{1\leq i,j\leq N} \det\Big(\frac{1-(u_p^2 e^{-i\phi_{k_l}})^{M-n}}{1-u_p^2 e^{-i\phi_{k_l}}}\Big)_{1\leq p,l\leq N}.
\end{aligned}
\tag{49}
$$

The Binet–Cauchy formula enables to evaluate (49) as follows:

$$
\begin{aligned}
& \langle \Psi_N(\mathbf{v}) \mid \bar{\varPi}_n\, e^{-\beta \hat{H}_{XX}}\, \bar{\varPi}_n \mid \Psi_N(\mathbf{u}) \rangle \;= \\
& = \frac{1}{\mathcal{V}(\mathbf{u}^2)\mathcal{V}(\mathbf{v}^{-2})} \det\left(\sum_{k,l=0}^{M-n-1} F_{k;\,l}(\beta)\, \frac{u_i^{2l}}{v_j^{2k}}\right)_{1\leq i,j\leq N},
\end{aligned}
\tag{50}
$$

where $F_{k;\,l}(\beta)$ is defined by (46). Clearly, the relation (50) at $\beta = 0$ is reduced to (36). Expression for $\mathcal{T}(\boldsymbol{\theta}, n, \beta)$ (25) can straightforwardly be obtained from (50) at coinciding parameters $\mathbf{u} = \mathbf{v}$ (being solutions of the Bethe equations) with the help of $\mathcal{N}^2$ (33):

$$
\mathcal{T}(\boldsymbol{\theta}, n, \beta) \;=\; \frac{1}{(M+1)^N} \det\left(\sum_{k,l=0}^{M-n-1} F_{k;\,l}(\beta)\, e^{i(l\theta_i - k\theta_j)}\right)_{1\leq i,j\leq N}.
$$

3 Strongly anisotropic XXZ chain

3.1 The Bethe state vectors and their form-factors

Let us turn to the strong anisotropy limit, $\Delta \to -\infty$, which is described by the Hamiltonian $\hat{H}_{\rm SA}$ (15). The corresponding state-vector is given by Eq. (3),

$$
\mid \Psi_N(\mathbf{u}) \rangle = \sum_{\widetilde{\boldsymbol{\lambda}} \subseteq \{(M-2(N-1))^N\}} S_{\widetilde{\boldsymbol{\lambda}}}(\mathbf{u}^2) \prod_{k=0}^{M} (\sigma_k^-)^{e_k} \mid \Uparrow \rangle\,, \tag{51}
$$

where $S_{\widetilde{\boldsymbol{\lambda}}}(\mathbf{u}^2)$ is given by (24), and summation goes over all non-strict partitions. Summation over strict partitions $\boldsymbol{\mu}$ with the elements respecting the condition $\mu_i > \mu_{i+1}+1$ is equivalent to that over the non-strict partitions $\widetilde{\boldsymbol{\lambda}}$, where $\widetilde{\boldsymbol{\lambda}} = \boldsymbol{\mu} - 2\boldsymbol{\delta}$ and $M + 2(1-N) \geq \widetilde{\lambda}_1 \geq \widetilde{\lambda}_2 \geq \cdots \geq \widetilde{\lambda}_N \geq 0$. The scalar product of the state-vector (51) to its conjugated (defined similarly to (28)) is given by the relation looking, in turn, similarly to (29):

$$\langle \Psi_N(\mathbf{v}) \,|\, \Psi_N(\mathbf{u})\rangle \,=\, \sum_{\widetilde{\boldsymbol{\lambda}} \subseteq \{(M-2(N-1))^N\}} S_{\widetilde{\boldsymbol{\lambda}}}(\mathbf{v}^{-2}) S_{\widetilde{\boldsymbol{\lambda}}}(\mathbf{u}^2) \,=\, \frac{\det(T_{kj})_{1\leq k,j\leq N}}{\mathcal{V}(\mathbf{u}^2)\mathcal{V}(\mathbf{v}^{-2})}\,, \tag{52}$$

where the entries T_{kj} take the form:

$$T_{kj} = \frac{1-(u_k^2/v_j^2)^{M-N+2}}{1-u_k^2/v_j^2}\,, \tag{53}$$

and the notation (30) for the Vandermonde determinant is used. When the sets of parameters $\mathbf{v}$ and $\mathbf{u}$ in left-hand side of (52) coincide, usage of the Bethe equations (18) enables to express the entries (53) as follows: $T_{jk} = 1 + (M-N+1)\delta_{jk}$. Therefore the norm of the Bethe eigen-vectors $\mathcal{N}^2(\boldsymbol{\theta}) \equiv \langle \Psi_N(\boldsymbol{\theta}) \mid \Psi_N(\boldsymbol{\theta})\rangle$ is given by

$$\mathcal{N}^2(\boldsymbol{\theta}) \,=\, \frac{(M+1)(M+1-N)^{N-1}}{\mathcal{V}(e^{i\boldsymbol{\theta}})\mathcal{V}(e^{-i\boldsymbol{\theta}})} \,=\, \frac{(M+1)(M+1-N)^{N-1}}{\prod\limits_{1\leq m<l\leq N} 2(1-\cos(\theta_l-\theta_m))}\,. \tag{54}$$

The exponential parametrization (19) is meant in (54) in the compact form (32). It can be shown that the scalar product $\langle \Psi_N(\mathbf{v}) \,|\, \Psi_N(\mathbf{u})\rangle$ (52) vanishes (i.e., the state-vectors are orthogonal) provided the parameters $\mathbf{u}$ and $\mathbf{v}$ are independent Bethe solutions.

The nominator of the ratio (34) is calculated in the same way as in the Section 2.1, i.e., by means of the Binet-Cauchy formula:

$$\begin{aligned}\langle \Psi_N(\mathbf{v}) \mid \bar{\Pi}_n \,|\Psi_N(\mathbf{u})\rangle \,&=\, \sum_{\widetilde{\boldsymbol{\lambda}} \subseteq \{(M-2N-n+1)^N\}} S_{\widetilde{\boldsymbol{\lambda}}}(\mathbf{v}^{-2}) S_{\widetilde{\boldsymbol{\lambda}}}(\mathbf{u}^2) \\ &= \frac{1}{\mathcal{V}(\mathbf{u}^2)\mathcal{V}(\mathbf{v}^{-2})} \det\Big(\frac{1-(u_k^2/v_j^2)^{M-N-n+1}}{1-u_k^2/v_j^2}\Big)_{1\leq j,k\leq N}\,.\end{aligned} \tag{55}$$

After use of (54) and (55), the answer for (34), taken on the solutions of the Bethe equation (19) (i.e., the Emptiness Formation Probability), appears in the following form:

$$\begin{aligned}\mathcal{T}(\boldsymbol{\theta}, n) = \frac{M-N+1}{M+1} \times \det\Big(&(1-\frac{n}{M-N+1})\delta_{jk} + \\ &+ \frac{1-e^{in(\theta_j-\theta_k)}}{(M-N+1)(1-e^{i(\theta_k-\theta_j)})}(1-\delta_{jk})\Big)_{1\leq k,j\leq N}\,.\end{aligned} \tag{56}$$

3.2 Thermal correlator of ferromagnetic string

Let us turn to obtaining of the average $\mathcal{T}(\mathbf{v}, \mathbf{u}, n, \beta)$ (38), where the Hamiltonian $\hat{H}_{XX}$ is replaced by $\hat{H}_{\mathrm{SA}}$ (15). Using (51) we obtain the corresponding nominator of (38):

$$\begin{aligned}&\langle \Psi_N(\mathbf{v}) \mid \bar{\Pi}_n \, e^{-\beta \hat{H}_{\mathrm{SA}}} \, \bar{\Pi}_n \,|\Psi_N(\mathbf{u})\rangle \,= \\ &= \sum_{\widetilde{\boldsymbol{\lambda}}^L,\, \widetilde{\boldsymbol{\lambda}}^R \subseteq \{(M-2N-n+1)^N\}} S_{\widetilde{\boldsymbol{\lambda}}^L}(\mathbf{v}^{-2})\, S_{\widetilde{\boldsymbol{\lambda}}^R}(\mathbf{u}^2)\, F_{\widetilde{\boldsymbol{\mu}}^L;\, \widetilde{\boldsymbol{\mu}}^R}(\beta)\,,\end{aligned} \tag{57}$$

where $F_{\widetilde{\boldsymbol{\mu}}^L;\,\widetilde{\boldsymbol{\mu}}^R}(\beta)$ is $2N$-point correlation function over the ferromagnetic state given, practically, by (40) excepting that $\hat{H}_{XX}$ (8) is replaced by $\hat{H}_{\rm SA}$ (15). Summations in (57) go over non-strict partitions $\widetilde{\boldsymbol{\lambda}}^L$ and $\widetilde{\boldsymbol{\lambda}}^R$ of the same kind like in (52). The corresponding strict partitions $\widetilde{\boldsymbol{\mu}}^L$ and $\widetilde{\boldsymbol{\mu}}^R$ are defined as follows: $\widetilde{\boldsymbol{\mu}}^{L,R} = \widetilde{\boldsymbol{\lambda}}^{L,R} + 2\boldsymbol{\delta}$, where $\boldsymbol{\delta}$ implies the partition $(N-1, N-2, \dots, 0)$. It is crucial that now the lattice indices $\widetilde{\boldsymbol{\mu}}^L$, $\widetilde{\boldsymbol{\mu}}^R$ respect the exclusion requirement: occupation of nearest sites is forbidden. Besides, an analogue of the relation (42) can be written as follows:

$$\begin{aligned}
&\mathcal{D}^K_{\beta/2}\Big[\langle\Psi_N(\mathbf{v}) \mid \bar{\Pi}_n\, e^{-\beta\hat{H}_{\rm SA}}\, \bar{\Pi}_n \mid \Psi_N(\mathbf{u})\rangle\Big] = \\
&= \sum_{\widetilde{\boldsymbol{\lambda}}^L,\, \widetilde{\boldsymbol{\lambda}}^R \subseteq \{(M-2N-n+1)^N\}} |P_K(\widetilde{\boldsymbol{\mu}}^R \to \widetilde{\boldsymbol{\mu}}^L)|\, S_{\widetilde{\boldsymbol{\lambda}}^L}(\mathbf{v}^{-2})\, S_{\widetilde{\boldsymbol{\lambda}}^R}(\mathbf{u}^2)\,.
\end{aligned} \tag{58}$$

The solutions of the Bethe equation (18) constitute a complete set of the eigen-states [53]. Taking into account the orthogonality of the corresponding states, one can consider the resolution of the identity operator:

$$\mathbb{I} = \sum_{\{\boldsymbol{\theta}\}} \mathcal{N}^{-2}(\boldsymbol{\theta})\, \mid \Psi_N(\boldsymbol{\theta})\rangle\langle\Psi_N(\boldsymbol{\theta}) \mid, \tag{59}$$

where summation goes over all independent solutions of the Bethe equation (18), and the square of the norm $\mathcal{N}^2(\boldsymbol{\theta})$ is given by (54). We shall calculate (57) inserting (59) into left-hand side of (57) and using appropriately Eq. (55). We take into account that

$$\langle\Psi_N(\mathbf{v}) \mid e^{-\beta\hat{H}_{\rm SA}} \mid \Psi_N(\boldsymbol{\theta})\rangle = \langle\Psi_N(\mathbf{v}) \mid \Psi_N(\boldsymbol{\theta})\rangle\, e^{-\beta E_N(\boldsymbol{\theta})},$$

where the expression for the energy $E_N(\boldsymbol{\theta})$ is given by the relation (21). Further, we use (55) and obtain:

$$\begin{aligned}
&\langle\Psi_N(\mathbf{v}) \mid \bar{\Pi}_n\, e^{-\beta\hat{H}_{\rm SA}}\, \bar{\Pi}_n \mid \Psi_N(\mathbf{u})\rangle \\
=\;& \frac{1}{(M+1)(M+1-N)^{N-1}\,\mathcal{V}(\mathbf{u}^2)\mathcal{V}(\mathbf{v}^{-2})} \sum_{M-N\ge l_1 > l_2 \dots > l_N \ge 0} e^{-\beta E_N(\boldsymbol{\theta})} \\
\times\;& \det\Big(\frac{1-(e^{i\theta_i}v_j^{-2})^{M-N-n+1}}{1-e^{i\theta_i}v_j^{-2}}\Big)_{1\le i,j\le N} \det\Big(\frac{1-(u_l^2 e^{-i\theta_p})^{M-N-n+1}}{1-u_l^2 e^{-i\theta_p}}\Big)_{1\le l,p\le N},
\end{aligned} \tag{60}$$

where summation goes over the ordered sets $\{I_k\}_{1\le k\le N}$, which parametrize the solution (19). Expression for $\mathcal{T}(\boldsymbol{\theta}, n, \beta)$ (25) can be obtained by means of (54) and (60), where it is necessary to put $\mathbf{u}^2 = \mathbf{v}^2 = e^{i\boldsymbol{\theta}}$, while $\boldsymbol{\theta}$ is the solution (20) of the Bethe equation for the ground state.

4 Boxed plane partitions

We shall show that the scalar products of the state vectors and the emptiness formation probability are related to the generating functions of the boxed plane partitions.

An array $(\pi_{i,j})_{i,j\ge 1}$ of non-negative integers that are non-increasing as functions of both i and j ($i, j \in \{1, 2, \dots\}$ is called a plane partition $\boldsymbol{\pi}$ [56]. The integers $\pi_{i,j}$ are called the parts of the plane partition, and $|\boldsymbol{\pi}| = \sum\limits_{i,j\ge 1} \pi_{i,j}$ is its volume. Each plane partition has a three dimensional diagram which can be interpreted as a stacks of unit cubes (three-dimensional Young diagram). The height of a stack with coordinates (i, j) is equal to the part of plane partition $\pi_{i,j}$. If we have $i \leq r$, $j \leq s$ and $\pi_{i,j} \leq t$ for all cubes of the plane partition, it is said that the plane partition is contained in a box with side lengths r, s, t. If $\pi_{i,j} > \pi_{i+1,j}$, i.e. if the parts of plane partition $\boldsymbol{\pi}$ are decaying along each column, then $\boldsymbol{\pi}$ is called column strict plane partition. We shall call the partition $\boldsymbol{\pi}$ that are decaying along each column and each raw ($\pi_{i,j} > \pi_{i+1,j}$ and $\pi_{i,j} > \pi_{i,j+1}$) as the strict plane partition. The element $\pi_{1,1}$ of the strict plane partition $\boldsymbol{\pi}$ satisfies the condition $\pi_{1,1} \geq 2r - 2$, if all $i, j \leq r$.

An arbitrary plane partition in a box $r \times r \times t$ may be transferred into a column strict plane partition in a box $r \times r \times (t + r - 1)$ by adding to an array $(\pi_{i,j})_{i,j \geq 1}$ the $r \times r$ matrix

$$\boldsymbol{\pi}_{\text{cspp}} = \begin{pmatrix} r-1 & r-1 & \cdots & r-1 \\ r-2 & r-2 & \cdots & r-2 \\ \vdots & \vdots & & \vdots \\ 0 & 0 & \cdots & 0 \end{pmatrix},$$

which corresponds to a minimal column strict plane partition. The volumes of the column strict plane partition and correspondent plane partition are related

$$|\boldsymbol{\pi}_{\text{cspp}}| = |\boldsymbol{\pi}| + \frac{N^2(N-1)}{2}. \tag{61}$$

An arbitrary plane partition in a box $r \times r \times t$ may be transferred into a strict plane partition in a box $r \times r \times (t + 2r - 2)$ by adding to an array $(\pi_{i,j})_{i,j \geq 1}$ the $r \times r$ matrix

$$\boldsymbol{\pi}_{\text{spp}} = \begin{pmatrix} 2r-2 & 2r-3 & \cdots & r-1 \\ 2r-3 & 2r-4 & \cdots & r-2 \\ \vdots & \vdots & & \vdots \\ r-1 & r-2 & \cdots & 0 \end{pmatrix},$$

which corresponds to a minimal strict plane partition. The volumes of the strict plane partition and correspondent plane partition are related

$$|\boldsymbol{\pi}_{\text{spp}}| = |\boldsymbol{\pi}| + N^2(N-1). \tag{62}$$

The partition function of the three dimensional Young diagrams, or saying differently the generating function of plane partitions is equal to

$$Z(q) = \sum_{\{\boldsymbol{\pi}\}} q^{|\boldsymbol{\pi}|}, \tag{63}$$

where q is a weight, and summation is performed over all plane partitions in a box. Formulas (61) and (62) provide the connection between partition functions of the plane partitions of different types. The generating functions of the column strict and strict plane partitions placed into a box $N \times N \times M$ are equal respectively to

$$Z_{\text{cspp}}(q) = q^{\frac{N^2}{2}(N-1)} \prod_{1 \leq j,k \leq N} \frac{1 - q^{M+1+j-k}}{1 - q^{j+k-1}}, \tag{64}$$

$$Z_{\text{spp}}(q) = q^{N^2(N-1)} \prod_{1 \leq j,k \leq N} \frac{1 - q^{M+3-j-k}}{1 - q^{j+k-1}}. \tag{65}$$

The scalar product (29) is related to the partition function of the column strict three-dimensional Young diagrams placed into $N \times N \times M$ box. Really, the parametrizations $v_j = q^{-\frac{j}{2}}$ and $u_j = q^{\frac{j-1}{2}}$ give:

$$\begin{aligned} &\langle \Psi_N(q^{-\frac{1}{2}}, \dots, q^{-\frac{N}{2}}) \,|\, \Psi_N(1, \dots, q^{\frac{N-1}{2}}) \rangle \;= \\ &= \sum_{\boldsymbol{\lambda} \subseteq \{(M+1-N)^N\}} S_{\boldsymbol{\lambda}}(q, \dots, q^N)\, S_{\boldsymbol{\lambda}}(1, \dots, q^{N-1}) \;= \\ &= \frac{1}{\mathcal{V}(q, \dots, q^N) \mathcal{V}(1, \dots, q^{N-1})} \det \Bigl(\frac{1 - s^{j+k-1}}{1 - q^{j+k-1}} \Bigr)_{1 \le j,k \le N}, \end{aligned} \tag{66}$$

where $s = q^{M+1}$, and

$$\mathcal{V}^{-1}(q, \dots, q^N)\, \mathcal{V}^{-1}(1, \dots, q^{N-1}) \;=\; q^{-\frac{N}{6}(N-1)(2N-1)} \prod_{1 \le k < j \le N} \bigl(1 - q^{j-k}\bigr)^{-2}. \tag{67}$$

The determinant in (66) was calculated in the paper [61] in connection with the alternating sign matrices enumeration problem:

$$\begin{aligned} &\det \Bigl(\frac{1 - s^{j+k-1}}{1 - q^{j+k-1}} \Bigr)_{1 \le j,k \le N} = \\ &= q^{\frac{N}{6}(N-1)(2N-1)} \prod_{1 \le k < j \le N} \bigl(1 - q^{j-k}\bigr)^2 \prod_{k,j=1}^{N} \frac{1 - s q^{j-k}}{1 - q^{j+k-1}}\,. \end{aligned} \tag{68}$$

Taking into account (64), (67), and (68), we obtain for (66):

$$\langle \Psi_N(q^{-\frac{1}{2}}, \dots, q^{-\frac{N}{2}}) \,|\, \Psi_N(1, \dots, q^{\frac{N-1}{2}}) \rangle = q^{-\frac{N^2}{2}(N-1)} Z_{\mathrm{cspp}}(q)\,. \tag{69}$$

Thus, Eq. (69) reads that the scalar product of two state-vectors coincides at $q = 1$ with the number of column strict partitions in a box $N \times N \times M$, i.e., with $Z_{\mathrm{cspp}}(1)$.

The same parametrizations $v_j = q^{-\frac{j}{2}}$ and $u_j = q^{\frac{j-1}{2}}$ enable to express the scalar product (52) corresponding to the strong anisotropy limit. The same representation is valid though $s = q^{M-N+2}$ now, while the range of summation over $\widetilde{\boldsymbol{\lambda}}$ takes the form:

$$\widetilde{\boldsymbol{\lambda}} \subseteq \{(M - 2(N-1))^N\}\,.$$

Taking into account Eq. (68), we now obtain:

$$\langle \Psi_N(q^{-\frac{1}{2}}, \dots, q^{-\frac{N}{2}}) \,|\, \Psi_N(1, \dots, q^{\frac{N-1}{2}}) \rangle = q^{-N^2(N-1)} Z_{\mathrm{spp}}(q)\,,$$

where $Z_{\mathrm{spp}}(q)$ is the generating function of strict plane partitions (65). The corresponding value of the scalar product at $q = 1$ coincides with the number of strict plane partitions $Z_{\mathrm{spp}}(1)$.

Now lets turn to the expectation value of the ferromagnetic string (36). In the present parametrization, we obtain (with regard at (67) and (68)):

$$\begin{aligned} &\langle \Psi_N(q^{-\frac{1}{2}}, \dots, q^{-\frac{N}{2}}) \,|\, \bar{\Pi}_n \,|\, \Psi_N(1, \dots, q^{\frac{N-1}{2}}) \rangle \;= \\ &= \sum_{\boldsymbol{\lambda} \subseteq \{(M-N-n)^N\}} S_{\boldsymbol{\lambda}}(q, \dots, q^N)\, S_{\boldsymbol{\lambda}}(1, \dots, q^{N-1}) \;=\; \prod_{k,j=1}^{N} \frac{1 - s q^{j-k}}{1 - q^{j+k-1}}\,, \end{aligned} \tag{70}$$

where $s = q^{M-n}$, i.e, (70) differs from (66) in the sense that $n+1$ is subtracted from $M+1$. The box containing the plane partitions is now of smaller maximal height: $N \times N \times (M-n-1)$. The expectation value in left-hand side of (70)), being considered at $q \to 1$, "counts" the number of plane partitions of smaller height. The same is true for the case of strong anisotropy, and (55) takes the form:

$$\begin{gathered}\langle \Psi_N(q^{-\frac{1}{2}}, \dots, q^{-\frac{N}{2}}) \mid \bar{\mathit{\Pi}}_n \mid \Psi_N(1, \dots, q^{\frac{N-1}{2}}) \rangle \;= \\ = \sum_{\widetilde{\boldsymbol{\lambda}} \subseteq \{(M-2N-n+1)^N\}} S_{\widetilde{\boldsymbol{\lambda}}}(q, \dots, q^N)\, S_{\widetilde{\boldsymbol{\lambda}}}(1, \dots, q^{N-1}) \;=\; \prod_{k,j=1}^{N} \frac{1 - sq^{j-k}}{1 - q^{j+k-1}}\,,\end{gathered} \tag{71}$$

where $s = q^{M-N-n+1}$.

5 Low temperature asymptotics

Let us go over to the case of long enough XX-chain, i.e., $M \gg 1$ while N is moderate: $M \gg N \gg 1$. Now the correlator $F_{k;l}(\beta)$ is approximately given by the modified Bessel function instead of (46):

$$F_{k;l}(\beta) = I_{k-l}(\beta) = \frac{1}{2\pi} \int\limits_{-\pi}^{\pi} e^{\beta \cos\phi} e^{i(k-l)\phi} d\phi\,. \tag{72}$$

In the limit of small "temperature" ($1/\beta \to 0$) and for moderate values of $m \equiv |k-l|$, we use the known asymptotics of the Bessel function and obtain:

$$F_{k;l}(\beta) \simeq \frac{e^{\beta}}{\sqrt{2\pi\beta}} \Bigl(1 - \frac{4m^2 - 1}{8\beta} + \dots\Bigr)\,, \tag{73}$$

i.e., the power decay is governed by the critical exponent $\xi = -1/2$.

Since the summations can be replaced by the integrations at large enough M, we obtain from (39) and (47):

$$\begin{aligned}\langle \Psi_N(\mathbf{v}) \mid \bar{\mathit{\Pi}}_n\, e^{-\beta \hat{H}_{XX}}\, \bar{\mathit{\Pi}}_n \mid \Psi_N(\mathbf{u}) \rangle \;=\; & \frac{e^{\beta N}}{N!} \prod_{i=1}^{N} \Bigl(\int\limits_{-\pi}^{\pi} \frac{d\phi_i}{2\pi} \Bigr)\, e^{-\beta \sum\limits_{l=1}^{N} (1 - \cos\phi_l)} \\ & \times \mathcal{P}(\mathbf{v}^{-2}, e^{i\boldsymbol{\phi}}) \mathcal{P}(e^{-i\boldsymbol{\phi}}, \mathbf{u}^2) \prod_{1 \le k < l \le N} |\, e^{i\phi_k} - e^{i\phi_l} \,|^2\,,\end{aligned} \tag{74}$$

where the continuous integration variables $\phi_i \in [0, 2\pi]$ are due to the change of the discrete variables $k_i \in \mathcal{M}$ as follows: $\phi_{k_i} \mapsto \phi_i$, $\forall i$. Further, the integral (74) can be approximated at β tending to infinity as follows:

$$\begin{aligned}\langle \Psi_N(\mathbf{v}) \mid \bar{\mathit{\Pi}}_n\, e^{-\beta \hat{H}_{XX}}\, \bar{\mathit{\Pi}}_n \mid \Psi_N(\mathbf{u}) \rangle \simeq \mathcal{P}(\mathbf{v}^{-2}, \mathbf{1}) \mathcal{P}(\mathbf{1}, \mathbf{u}^2)\, \frac{e^{\beta N}}{(2\pi)^N N!} \\ \times \int\limits_{-\infty}^{\infty} \int\limits_{-\infty}^{\infty} \cdots \int\limits_{-\infty}^{\infty} e^{-(\beta/2) \sum\limits_{l=1}^{N} \phi_l^2} \prod_{1 \le k < l \le N} |\, \phi_k - \phi_l \,|^2\, d\phi_1 d\phi_2 \dots d\phi_N\,.\end{aligned} \tag{75}$$

The bold-faced argument $\mathbf{1}$ in the sum, say, $\mathcal{P}(\mathbf{1}, \mathbf{u}^2)$ implies that $S_{\boldsymbol{\lambda}}(e^{\pm i\boldsymbol{\phi}})$ in (48) is replaced by $S_{\boldsymbol{\lambda}}(1, 1, \dots, 1)$ provided the N-tuple of the exponentials $e^{\pm i\boldsymbol{\phi}}$ is substituted by N-tuple of unities. The corresponding value of the Schur function can be obtained [56]:

$$S_{\boldsymbol{\lambda}}(1, 1, \dots, 1) = \frac{\prod\limits_{1\le k<l\le N} (\lambda_l - l - \lambda_k + k)}{(N-1)!(N-2)! \cdots 1!0!} \,. \tag{76}$$

Right-hand side of (76) coincides with the dimensionality $\mathsf{d}_{\boldsymbol{\lambda}} \equiv \dim \boldsymbol{\pi}_{\boldsymbol{\lambda}}$ of the unitary irreducible representations of the unitary group $\mathcal{U}(N)$, which corresponds to the signature $\boldsymbol{\lambda}$, i.e., $\mathsf{d}_{\boldsymbol{\lambda}} = S_{\boldsymbol{\lambda}}(1, 1, \dots, 1)$ [58]. The Schur functions provide a base of the ring of symmetric polynomials of N variables [56].

The integral in (75) is the *Mehta integral* [59] of the *Gaussian Unitary Ensemble* of random matrices. Its value is known, and the estimate for right-hand side of (75) acquires the form:

$$\langle \Psi_N(\mathbf{v}) \mid \bar{\varPi}_n \, e^{-\beta \hat{H}_{XX}} \, \bar{\varPi}_n \mid \Psi_N(\mathbf{u}) \rangle \simeq \mathcal{P}(\mathbf{v}^{-2}, \mathbf{1}) \, \mathcal{P}(\mathbf{1}, \mathbf{u}^2) \, \frac{e^{\beta N}}{\beta^{N^2/2}} \prod_{n=1}^{N} \frac{\Gamma(n)}{(2\pi)^{1/2}} \,. \tag{77}$$

Let us consider the case when $\mathbf{u}^2$ and $\mathbf{v}^2$ in (77) are the solutions of the Bethe equations (10). Approximate estimates $\mathbf{u}^2 = e^{i\boldsymbol{\theta}} \simeq \mathbf{1}$ and $\mathbf{v}^{-2} = e^{-i\boldsymbol{\theta}} \simeq \mathbf{1}$ are valid at $1 \ll N \ll M$. Thus, the estimate (77) takes the limiting form:

$$\begin{aligned} &\langle \Psi_N(1, \dots, 1) \mid \bar{\varPi}_n \, e^{-\beta \hat{H}_{XX}} \, \bar{\varPi}_n \mid \Psi_N(1, \dots, 1) \rangle \simeq \\ &\simeq \Big(\sum_{\boldsymbol{\lambda} \subseteq \{(M-N-n)^N\}} \mathsf{d}_{\boldsymbol{\lambda}} \, \mathsf{d}_{\boldsymbol{\lambda}} \Big)^2 \, \frac{e^{\beta N}}{\beta^{N^2/2}} \prod_{n=1}^{N} \frac{\Gamma(n)}{(2\pi)^{1/2}} \,. \end{aligned} \tag{78}$$

The present estimate is proportional to the square of (70) at $q = 1$, i.e., to the square of the number of boxed plane partitions of the size $N \times N \times (M - n - 1)$. As a result, the ratio of the average $\mathcal{T}(\boldsymbol{\theta}, n, \beta)$ (25) to the partiton function of the XX model $\mathcal{Z}$ [39] can be estimated as follows:

$$\frac{\mathcal{T}(\boldsymbol{\theta}, n, \beta)}{\mathcal{Z}} \simeq \text{const} \times \Big(\prod_{k,j=1}^{N} \frac{M - n + j - k}{j + k - 1} \Big)^2 \, \frac{e^{-\beta \mathcal{E}_0^{XX}}}{\beta^{N^2/2}} \,,$$

where $\mathcal{E}_0^{XX}$ is the ground state energy of the XX model. In the same limit, the generating function (42) is specified as follows:

$$\begin{aligned} &\mathcal{D}^K_{\beta/2} \Big[\langle \Psi_N(1, \dots, 1) \mid \bar{\varPi}_n \, e^{-\beta \hat{H}_{XX}} \, \bar{\varPi}_n \mid \Psi_N(1, \dots, 1) \rangle \Big] = \\ &= \sum_{\boldsymbol{\lambda}^L, \boldsymbol{\lambda}^R \subseteq \{(M-N-n)^N\}} |P_K(\boldsymbol{\mu}^R \to \boldsymbol{\mu}^L)| \, \mathsf{d}_{\boldsymbol{\lambda}^L} \, \mathsf{d}_{\boldsymbol{\lambda}^R} \,, \end{aligned}$$

where a relationship between strict and non-strict partitions is valid: $\boldsymbol{\mu}^{L,R} = \boldsymbol{\lambda}^{L,R} + \boldsymbol{\delta}$ (see, for instance, (39)).

In the case of strong anisotropy, we use (71) and obtain an analogous estimate which demonstrates the proportionality to the square of the number of strict plane partitions in a box of the size $N \times N \times (M - n - N)$:

$$\begin{aligned}
\frac{\mathcal{T}(\boldsymbol{\theta}, n, \beta)}{\mathcal{Z}} \; &\simeq \; \text{const} \times \Bigl(\sum_{\widetilde{\boldsymbol{\lambda}} \subseteq \{(M-2N-n+1)^N\}} \mathsf{d}_{\widetilde{\boldsymbol{\lambda}}}\, \mathsf{d}_{\widetilde{\boldsymbol{\lambda}}} \Bigr)^2 \frac{e^{-\beta \mathcal{E}_0^{\mathrm{SA}}}}{\beta^{N^2/2}} \; = \\
&= \; \text{const} \times \Bigl(\prod_{k,j=1}^{N} \frac{M - N + 1 - n + j - k}{j + k - 1} \Bigr)^2 \frac{e^{-\beta \mathcal{E}_0^{\mathrm{SA}}}}{\beta^{N^2/2}} \, ,
\end{aligned}$$

where $\mathcal{Z}$ is the partition function of the model in the limit of strong anisotropy [5, 8], and $\mathcal{E}_0^{\mathrm{SA}}$ is the corresponding ground state energy. The relation (58) is specified as follows:

$$\begin{aligned}
&\mathcal{D}^{K}_{\beta/2} \Bigl[\langle \Psi_N(1, \dots, 1) \mid \bar{\mathit{\Pi}}_n \, e^{-\beta \hat{H}_{\mathrm{SA}}} \, \bar{\mathit{\Pi}}_n \mid \Psi_N(1, \dots, 1) \rangle \Bigr] \; = \\
&= \sum_{\widetilde{\boldsymbol{\lambda}}^L, \widetilde{\boldsymbol{\lambda}}^R \subseteq \{(M-2N-n+1)^N\}} |P_K(\widetilde{\boldsymbol{\mu}}^R \to \widetilde{\boldsymbol{\mu}}^L)| \, \mathsf{d}_{\widetilde{\boldsymbol{\lambda}}^L} \, \mathsf{d}_{\widetilde{\boldsymbol{\lambda}}^R} \, ,
\end{aligned}$$

where $\widetilde{\boldsymbol{\mu}}^{L,R} = \widetilde{\boldsymbol{\lambda}}^{L,R} + 2\boldsymbol{\delta}$ (see (57)).

6 Discussion

The XXZ Heisenberg chain has been considered for two specific limits of the anisotropy parameter: $\Delta \to 0$ and $\Delta \to -\infty$. The corresponding state-vectors have been expressed by means of the symmetric Schur functions. Certain expectation values and thermal correlation functions of the ferromagnetic string operators have been calculated over the base of N-particle Bethe states. The expectation values obtained are of the type of the emptiness formation probability. The thermal correlator of the ferromagnetic string operator is expressed through the generating function of the lattice paths of random walks of vicious walkers. The thermal correlator in question turns out to be a generating function of certain polynomials build up from the Schur polynomials. A relationship between the expectation values obtained and the generating functions of boxed plane partitions is discussed. Asymptotic estimate of the thermal expectation value of the ferromagnetic string is obtained in the limit of zero temperature for $\Delta = 0$. These estimates are expressed in terms of the dimensionality of the irreducible representations of the group $\mathcal{U}(N)$.

References

[1] W. Heisenberg, *Zur Theorie des Ferromagnetismus*, Zeitschrift für Physik, **49** (1928), No. 9-10, 619–636.

[2] H. Bethe, *Zur Theorie der Metalle. I. Eigenwerte und Eigenfunktionen Atomkette*, Zeitschrift für Physik, **71** (1931), No. 3-4, 205–226.

[3] C. N. Yang, C. P. Yang, *One-dimensional chain of anisotropic spin-spin interactions. I. Proof of Bethe's hypothesis for ground state in a finite system*, Phys. Rev., **150** (1966), No. 1, 321–327.

[4] C. N. Yang, C. P. Yang, *One-dimensional chain of anisotropic spin-spin interactions. II. Properties of the ground state energy per lattice site for an infinite system*, Phys. Rev., **150** (1966), No. 1, 327–339.

[5] C. N. Yang, C. P. Yang, *One-dimensional chain of anisotropic spin-spin interactions. III. Applications*, Phys. Rev., **151** (1966), No. 1, 258–264.

[6] E. H. Lieb, F. Y. Wu, *Two dimensional ferroelectric models*, In: Phase transitions and critical phenomena, vol. 1, Eds., C. Domb, M. Green, Academic Press, London, 1972, pp. 331–490.

[7] R. J. Baxter, *Exactly Solved Models in Statistical Mechanics*, Academic Press, London, 1982.

[8] M. Gaudin, *La Fonction d'Onde de Bethe*, Masson, Paris, 1983.

[9] L. D. Faddeev, *Quantum completely integrable models of field theory*, Sov. Sci. Rev. Math. C, **1** (1980), 107–160; In: 40 Years in Mathematical Physics, World Sci. Ser. 20th Century Math., vol. 2, World Sci., Singapore, 1995, pp. 187–235.

[10] P. P. Kulish, E. K. Sklyanin, *Quantum spectral transform method. Recent developments*, Lecture Notes in Phys., vol. 151, Springer, Berlin, etc., 1982, pp. 61–119.

[11] L. D. Faddeev, L. A. Takhtajan, *Quantum inverse scattering method and the XYZ Heisenberg model*, Usp. Mat. Nauk, **34** (1979), No. 5(209), 13–63. [In Russian]

[12] L. D. Faddeev, L. A. Takhtajan, *What is the spin of a spin wave*, Phys. Lett. A, **85** (1981), No. 6-7, 375–377.

[13] V. E. Korepin, *Calculation of norms of Bethe wave functions*, Comm. Math. Phys. **86** (1982), No. 3, 391–418.

[14] A. G. Izergin, V. E. Korepin, *Correlation functions for the Heisenberg XXZ-antiferromagnet*, Comm. Math. Phys. **99** (1985), No. 2, 271–302.

[15] P. P. Kulish, F. A. Smirnov, *Anisotropic Heisenberg ferromagnet with a ground of the domain wall type*, J. Phys. C: Solid State Phys., **18** (1985), No. 5, 1037–1048.

[16] N. M. Bogoliubov, A. G. Izergin, V. E. Korepin, *Correlation Functions of Integrable Systems and the Quantum Inverse Scattering Method*, Nauka, Moscow, 1992. [In Russian]

[17] V. E. Korepin, N. M. Bogoliubov, A. G. Izergin, *Quantum Inverse Scattering Method and Correlation Functions*, Cambridge University Press, Cambridge, 1993.

[18] F. H. L. Eßler, H. Frahm, A. G. Izergin, V. E. Korepin, *Determinant representation for correlation functions of spin-1/2 XXX and XXZ Heisenberg magnets*, Comm. Math. Phys., **174** (1995), No. 1, 191–214.

[19] N. Kitanine, J. M. Maillet, V. Terras, *Form factors of the XXZ Heisenberg spin-$\frac{1}{2}$ finite chain*, Nucl. Phys. B **554** (1999), No. 3, 647–678.

[20] N. Kitanine, J. M. Maillet, N. Slavnov, V. Terras, *Correlation functions of the XXZ spin-$\frac{1}{2}$ Heisenberg chain at the free fermion point from their multiple integral representations*, Nucl. Phys. B **642** (2002), No. 3, 433–455.

[21] M. E. Fisher, *Walks, walls, wetting, and melting*, J. Statist. Phys., **34** (1984), No. 5–6, 667–729.

[22] P. J. Forrester, *Exact solution of the lock step model of vicious walkers*, J. Phys. A.: Math. Gen., **23** (1990), No. 7, 1259–1273.

[23] T. Nagao, P. J. Forrester, *Vicious random walkers and a discretization of Gaussian random matrix ensembles*, Nucl. Phys. B, **620** (2002), No. 3, 551–565.

[24] A. J. Guttmann, A. L. Owczarek, X. G. Viennot, *Vicious walkers and Young tableaux I: without walls*, J. Phys. A: Math. Gen., **31** (1998), No. 40, 8123–8135.

[25] C. Krattenthaler, A. J. Guttmann, X. G. Viennot, *Vicious walkers, friendly walkers and Young tableaux: II. With a wall*, J. Phys. A: Math. Gen., **33** (2000), No. 48, 8835–8866.

[26] C. Krattenthaler, A. J. Guttmann, X. G. Viennot, *Vicious walkers, friendly walkers, and Young tableaux. III. Between two walls*, J. Statist. Phys., **110** (2003), No. 3–6, 1069–1086.

[27] M. Katori, H. Tanemura, *Scaling limit of vicious walks and two-matrix model*, Phys. Rev. E, **66** (2002), No. 1, 011105 [12 pages].

[28] M. Katori, H. Tanemura, T. Nagao, N. Komatsuda, *Vicious walks with a wall, noncolliding meanders, and chiral and Bogoliubov–de Gennes random matrices*, Phys. Rev. E, **68** (2003), No. 2, 021112 [16 pages].

[29] M. Katori, H. Tanemura, *Nonintersecting paths, noncolliding diffusion processes and representation theory*, In: Combinatorial Methods in Representation Theory and their Applications. RIMS Kokyuroku, **1438** (2005), 83–102.

[30] P. Bak, C. Tang, K. Wiesenfeld, *Self-organized criticality*, Phys. Rev. A, **38** (1998), No. 1, 364–374.

[31] D. Huse, M. Fisher, *Commensurate melting, domain walls, and dislocations*, Phys. Rev. B, **29** (1984), No. 1, 239–270.

[32] J. W. Essam, A. J. Guttmann, *Vicious walkers and directed polymer networks in general dimensions*, Phys. Rev. E, **52** (1995), No. 6, 5849–5862.

[33] S. Yu. Grigoriev, V. B. Priezzhev, *Random walk of annihilating particles on the ring*, Teoret. Mat. Fiz., **146** (2006), No. 3, 488–498. [In Russian]

[34] J. W. van de Leur, A. Yu. Orlov, *Random turn walk on a half line with creation of particles at the origin.* arXiv:0801.0066

[35] N. M. Bogoliubov, *XX Heisenberg chain and random walks*, Zap. Nauchn. Sem. POMI **325** (2005), 13–27; English transl., J. Math. Sci. **138** (2006), No. 3, 5636–5643.

[36] N. M. Bogoliubov, *The integrable models for the vicious and friendly walkers*, Zap. Nauchn. Sem. POMI **335** (2006), 59–74; English transl., J. Math. Sci., **143** (2007), No. 1, 2729–2737.

[37] N. M. Bogoliubov, C. Malyshev, *A path integration approach to the correlators of XY Heisenberg magnet and random walks*, In: Proceedings of the 9th Intern. Conf. “Path Integrals: New Trends and Perspectives” (Dresden, Germany, September 23–28, 2007). Eds., W. Janke, A. Pelster (World Sci., Singapore, 2008), pp. 508–513. arXiv:0810.4816

[38] N. M. Bogoliubov, C. Malyshev, *The correlation functions of the XX Heisenberg magnet and random walks of vicious walkers*, Theor. Math. Phys., **159** (2009), No. 2, 563–574. arXiv:0903.3227

[39] E. Lieb, T. Schultz, D. Mattis, *Two soluble models of an antiferromagnetic chain*, Ann. Phys. (NY), **16** (1961), No. 3, 407–466.

[40] Th. Niemeijer, *Some exact calculations on a chain of spin* $\frac{1}{2}$*. I, II*, Physica, **36** (1967), No. 3, 377–419; **39** (1968), No. 3, 313–326.

[41] F. Colomo, A. G. Izergin, V. E. Korepin, V. Tognetti, *Correlators in the Heisenberg XXO chain as Fredholm determinants*, Phys. Lett. A, **169** (1992), No.4, 243–247.

[42] F. Colomo, A. G. Izergin, V. E. Korepin, V. Tognetti, *Temperature correlation functions in the XXO Heisenberg chain*, Theor. Math. Phys., **94** (1993), No.1, 11–38.

[43] A. G. Izergin, N. A. Kitanin, N. A. Slavnov, *On correlation functions of the XY model*, Zap. Nauchn. Semin. POMI, **224** (1995), 178–191; English transl., J. Math. Sci., **88** (1998), No.2, 224–232.

[44] A. G. Izergin, V. S. Kapitonov, N. A. Kitanin, *Equal-time temperature correlators of the one-dimensional Heisenberg XY chain*, Zap. Nauchn. Semin. POMI, **245** (1997), 173–206; English transl., J. Math. Sci., **100** (2000), No.2, 2120–2140.

[45] V. S. Kapitonov, A. G. Pronko, *Time-dependent temperature correlators of local spins of the one-dimensional Heisenberg XY chain*, Zap. Nauchn. Semin. POMI, **269** (2000), 216–261; English transl., J. Math. Sci., **115** (2003), No. 1, 2009–2032.

[46] C. Malyshev, *Functional integration with an “automorphic” boundary condition and correlators of third components of spins in the XX Heisenberg model*, Theor. Math. Phys., **136** (2003), No. 2, 1143–1154. arXiv:hep-th/0204007

[47] C. Malyshev, *Functional integration with “automorphic” boundary conditions and correlators of z-components of spins in the XY and XX Heisenberg chains*, In: *New Developments in Mathematical Physics Research*, Ed. Charles V. Benton (Nova Science Publishers, New York, 2004), pp. 85–116. arXiv:math-ph/0405009

[48] S. Sachdev, *Quantum Phase Transitions*, Cambridge University Press, Cambridge, 1999.

[49] F. Colomo, A. G. Izergin, V. Tognetti, *Correlation functions in the XXO Heisenberg chain and their relations with spectral shapes*, J. Phys. A: Math. Gen., **30** (1997), No. 2, 361–370.

[50] V. Korepin, J. Terilla, *Thermodynamic interpretation of quantum error correcting criterion*, Quantum Information Processing, **1** (2002), No. 4, 225–242.

[51] B.-Q. Jin, V. E. Korepin, *Entanglement, Toeplitz determinants and Fisher-Hartwig conjecture*, J. Stat. Phys., **116** (2004), No. 1-4, 79–95.

[52] F. C. Alcaraz, R. Z. Bariev, *An exactly solvable constrained XXZ chain*, In: *Statistical Physics on the Eve of the 21st Century. In Honour of J. B. McGuire on the Occasion of His 65th Birthday (Series on Advances in Statistical Mechanics)*, Eds., M. T. Batchelor, L. T. Wille (World Scientific, Singapore, 1999). arXiv:cond-mat/9904042

[53] N. I. Abarenkova, A. G. Pronko, *The temperature correlator in the absolutely anisotropic Heisenberg XXZ-magnet*, Theor. Math. Phys., **131** (2002), No. 2, 690–703.

[54] A. J. A. James, W. D. Goetze, F. H. L. Essler, *Finite-temperature dynamical structure factor of the Heisenberg-Ising chain*, Phys. Rev. B, **79** (2009), No. 21, 214408 [20 pages].

[55] P. Lu, G. Muller, M. Karbach, *Quasiparticles in the XXZ model.* arXiv:0909.2728

[56] I. G. Macdonald, *Symmetric Functions and Hall Polynomials*, Oxford University Press, Oxford, 1995.

[57] F. R. Gantmakher, *Theory of Matrices*, Nauka, Moscow, 1988. [In Russian]

[58] D. P. Zhelobenko, A. I. Stern, *Representations of the Lie Groups*, Nauka, Moscow, 1983. [In Russian]

[59] M. L. Mehta, *Random Matrices*, Academic Press, London, 1991.

[60] A. Okounkov, *Infinite wedge and random partitions*, Selecta Mathematica, New Series, **7** (2001), No. 1, 57–81.

[61] G. Kuperberg, *Another proof of the alternative-sign matrix conjecture*, Int. Math. Res. Notices, **1996** (1996), 139–150.